

1  
**Contains No CBI**



**PHILLIPS PETROLEUM COMPANY**  
BARTLESVILLE, OKLAHOMA 74004 918 661-6600

**"Contains NO CBI"**

HEALTH, ENVIRONMENT AND SAFETY

A

August 24, 1992

Compliance Audit Program  
CAP ID#: 8ECAP-0075

1992 SEP -2 PM 1:16  
OTS CBIC

**CERTIFIED MAIL - RETURN RECEIPT**

Document Processing Center (TS-790)  
Office of Pollution Prevention and Toxics  
Environmental Protection Agency  
401 M Street, SW  
Washington, D. C. 20460

8EHQ-92-12578

88920010762

INIT

Attn: Section 8(e) Coordinator  
(CAP Agreement)

Gentlemen:

Phillips Petroleum Company is submitting the enclosed sixty (60) reports (two boxes, numbered 1 and 2) of toxicological studies pursuant to category II.B.2.b of the CAP Agreement 8ECAP-0075 Reports. Reports being submitted contain no confidential business information.

We are sending an additional five boxes (box numbers 3-7) of reports of studies that have, previously, been submitted to the FYI coordinator of the Office of Pollution Prevention and Toxics by the American Petroleum Institute (API). These are being provided solely for the Agency's convenience.

For questions concerning this correspondence, please contact Fred Marashi at 918-661-8153.

Very truly yours,

Barbara J. Price  
Vice President  
Health, Environment & Safety

Enclosure (Seven Boxes)

FFM/dh:29

3/7/95



**Phillips Petroleum Company**

CAP Identification Number: 8ECAP-0075  
Pursuant to Category: II.B.2.b

44

**Contains No CBI**

**Title of Study:** Chronic Benzene Toxicology

**Name of Chemical:** Benzene

**CAS#:** 71-43-2

**Summary:** Rodents exposed chronically to benzene exhibited lymphoma, leukemia and other blood changes.

---

**Contact:**

Fred Marashi  
Phillips Petroleum Company  
13 D2 PB  
Bartlesville, OK 74004  
Phone: 918/661-8153  
Fax: 918/661-5664

2321

FINAL

REPORT

To the American Petroleum Institute  
Washington, D.C.


CHRONIC BENZENE TOXICOLOGY

Project U-150-14 (PS-7)

Principal Investigators: Sidney Laskin and Bernard Goldstein

From the Institute of Environmental Medicine  
New York University Medical Center  
New York, NY 10016

August 22, 1978

  
Carroll A. Snyder Ph.D.  
Research Assistant Professor

### Introduction

The rationale of this study was to develop suitable animal models for benzene-induced hematopoietic toxicity. The study was essentially divided into three sections or phases. Phase I was a comprehensive literature review which served as a basis for the direction of research and which was summarized in a monograph. Phase II included the toxicity studies of inhaled benzene and was divided into three areas. Area A was concerned with the inhalation protocols and peripheral blood cell monitoring. Area B dealt with the evaluation of hematological responses other than peripheral cell counts. Area C concerned the effects of benzene exposure on proliferating blood cell precursors as reflected by changes in bone marrow cytogenetics and colony forming abilities. Phase III dealt with the study of the metabolism of benzene which was investigated through pharmacokinetic techniques.

### Phase I - Literature Review

Over 1000 pertinent articles were critically reviewed and catalogued. A monograph based on this review was published in November, 1977 as a supplement to the Journal of Toxicology and Environmental Health.<sup>1</sup>

## Phase II - Toxicity

### Area A

Exposures were performed in identical inhalation units (1.6m<sup>3</sup> total volume). Chamber atmospheres were sampled every 0.5 hours or 12 times during the daily runs. For most studies treated animals received either 300 ppm or 100 ppm benzene vapor for 6 hr/d x 5 d/wk x life. Air sham controls were exposed analogously to filtered, conditioned air. When not in the inhalation units rats were housed two to a cage and fed food and water ad libitum. Mice were housed five to a cage in polycarbonate boxes and allowed food and water ad libitum. Animals received neither food nor water during exposures. The numbers of animals used in each study and calculated median lifespans are shown in Table 1.

Animals were usually bled (by tail clipping) biweekly for determination of peripheral cell responses. Only males were used in order to avoid anomolous white counts due to estrus. Test and corresponding air sham animals were bled within one hour by the same individual in order to minimize differences in white counts due to handling, circadian rhythm, etc. Routinely, red blood counts, white blood counts and differential white blood counts were determined on all animals, but other parameters were frequently investigated including hematocrit, hemoglobin and reticulocyte levels.

Sprague-Dawley Rats at 300 ppm

The chronic exposure of Sprague-Dawley rats to 300 ppm benzene was completed at 99 weeks with the death of the last test animal. These animals received up to 463 exposures over the 691 days of the study at an integrated mean concentration of 301 ppm. The calculated median lifespan of the test animals was 59 weeks versus 73 weeks for the controls (Table 1). Test animals first showed weight depression when compared with controls at 30 weeks, and this trend continued throughout the study although weight differences were never greater than two standard errors ( $\pm 2S_x$ ).

The acquisition of blood data was suspended after one year of exposure due to advancing mortality in test rats. Blood counts performed during the first year demonstrated that benzene induced significant ( $\pm 2S_x$ ) leukopenia in test rats and that this leukopenia was due to a selective decrease in lymphocyte levels (Figure 1). Red cell counts for the first exposure year suggested a trend to anemia in test rats, however, differences in red cell levels between test and control animals were not as striking as differences in white cell levels (Table 2). An evaluation of the weekly red cell data shows that test animals had lower counts than controls in 69% (26/38) of the counts taken during the first exposure year. Consistent with the red cell counts, hematocrit and hemoglobin determinations showed no differences between test and control animals.

Evaluation of tissue sections revealed hemosiderin pigments in 42% of the spleens taken from test rats as opposed to 18% in the spleens of control animals. These results were shown not to be significant, however, at the  $p = 0.05$  level by  $\chi^2$  analysis. Fatty changes appeared in 77% of the bone marrows of test animals versus 42% for controls. Evaluation by  $\chi^2$  showed these results to be significant at  $p = 0.05$ . No pathological differences were observed in the other organs studied.

#### AKR Mice at 300 ppm

The chronic exposure of AKR/J mice to 300 ppm benzene was completed after 28 weeks with the death of the last test animal. Treated mice received up to 123 exposures over the 181 days of the study at an integrated mean benzene concentration of 301 ppm.

Treated mice exhibited a median survival time of 17 weeks compared with a median survival time of 47 weeks for controls (Table 1). Weight data taken during the exposure showed severe weight loss for the treated versus the untreated mice (Table 3).

The monitoring of blood parameters was suspended after 92 days because of high mortality in the test mice. The blood data demonstrated that benzene induced severe lymphocytopenia in the exposed animals (Figure 2) but that these animals still retained an ability to produce granulocytes

during the exposure. Red cell data showed that benzene also induced anemia in test mice (Figure 3). Reticulocytosis first appeared concurrent with the development of anemia and became significant ( $\pm 2S_x$ ) at 37 days exposure (Table 4).

Evidence of malignant lymphoma was found in tissue sectioned from 91% of the control mice but in only 2% of the exposed mice. The AKR strain spontaneously develops lymphoid leukemia at an incidence of 85-95%. The very low lymphoma incidence in treated mice was due to the severe rate of mortality induced by the exposure. The first lymphoma appeared after 6 months in the untreated mice whereas the median survival for treated mice was only 4 months.

Hemosiderin pigments were found in spleens taken from 25% of the treated mice while no pigment was detected in control mice. Chi square analysis showed these differences to be significant at the  $p < 0.001$  level.

Bone marrow hypoplasia was exhibited by 81% of the exposed mice but by only 6% of the controls. This, also, was found to be significant at the  $p < 0.001$  level by  $X^2$  analysis. Several test mice exhibited bone marrow hypoplasia of such severe proportions that only isolated nests of hematopoietic tissue were present in an otherwise empty marrow.

#### C-57 B1 Mice at 300 ppm

The chronic exposure to C-57 B1 mice was completed after 488 days with the death of the last test animal.



During the exposure period, test mice received 331 exposures at an integrated mean concentration of 300 ppm.

Treated mice exhibited a median survival time of 49 weeks versus 83 weeks for controls (Table 1) and showed weight gain depression relative to controls throughout the study.

Acquisition of peripheral blood data was suspended after 61 weeks due to advancing mortality in test mice.

Test mice exhibited statistically significant lymphocytopenia and anemia relative to controls after one exposure week and this trend continued throughout the course of the study (Table 5). Granulocytosis was evident in test mice vis a vis controls after 17 weeks and was statistically significant during most of the remainder of the exposure (Table 5).

Malignant lymphoma was detected in 9/40 treated animals but in only 2/40 controls. This is a statistically significant increase in the treated animals at the  $p < 0.05$  level as shown by  $\chi^2$  analysis. Because C-57 B1 mice carry a radiation-inducible lymphoma virus,<sup>2</sup> the observed increased incidence of lymphoma may indicate similar modes of action for benzene and ionizing radiation. Six of the test mice presented hypoplastic bone marrow. Four of these were associated with malignant lymphoma. No control mice gave evidence of bone marrow hypoplasia.

Of those test animals that did not show evidence of malignant lymphoma, 13/31 exhibited bone marrow hyperplasia and 15/31 exhibited spleen hyperplasia. These compared with no cases of bone marrow hyperplasia and 2/38 cases of spleen hyperplasia in the controls.

Charles River CD-1 Mice at 300 ppm

The exposure of CD-1 mice to 300 ppm benzene ended after 222 days with the development of two cases of myelogenous leukemia. During the exposure period animals received 149 exposures at an integrated mean benzene concentration of 300 ppm. At the time exposure was suspended, 9 test animals remained alive. The last of these animals died 8 weeks later.

The calculated median survival time was 31 weeks for treated animals and 50 weeks for controls (Table 1). Treated animals showed a decreased rate of weight gain relative to controls throughout the exposure but the largest differences were noted from the twenty-eighth week onward.

The blood data presented below was compiled from all test and control animals with the exception of the two treated animals with myelogenous leukemia.

Treated animals exhibited statistically significant lymphocytopenia and anemia after one exposure week and this trend continued throughout the course of the study (Figures 4 and 5). At 25 weeks, treated mice began showing a steadily

increasing granulocytosis relative to controls which continued until the end of the study (Figure 4).

Histological evaluation of all animals is not yet complete, however, a summary of the findings for three mice presenting hematological and histological profiles of note is given below.

A mouse dying after 130 exposure showed a white count in excess of  $160,000/\text{mm}^3$  (control= $12,000/\text{mm}^3$ ). The peripheral smear showed a preponderance of granulocytes shifted to the left. Bone marrow, spleen and liver sections were perfused with myeloid cells and there was infiltration by these cells into the muscle and connective tissue surrounding the bone. Death was ascribed to chronic myelogenous leukemia.

A second mouse dying after 139 exposures showed a normal white count of  $11,000/\text{mm}^3$  but 30% of the cells were blasts. Other cells in the peripheral smear were granulocytic types shifted to the left. Masses of cells were observed replacing normal architecture in bone marrow and spleen. These cells were a mononuclear type of varying sizes and appeared to mature towards the granulocytic series. Death was ascribed to acute myelogenous leukemia. Originally there was some doubt as to the nature of the proliferating cell type in this case. The peripheral cell morphology could have been consistent with either stem cell leukemia or acute myelogenous leukemia. Since there is a 4% spontaneous incidence of stem cell leukemia associated with these mice,<sup>3</sup>

the question of cell type as it concerns benzene exposure was crucial. Review of pertinent slides by Drs. Arthur Upton and Albert Jonas (see enclosures) seems to confirm the myeloid nature of the proliferating cell. Dr. Jonas' diagnosis is particularly compelling since he has, to our knowledge, reviewed the largest colony of these animals.

In addition to the above cases, a CD-1 mouse dying after 222 exposures showed alternating elevated and normal white counts during successive monitorings. The peripheral smears showed granulocytes shifted to the left. The bone marrow was perfused with granulocytic cells and contained few other myelopoietic cell types. There was no invasion by these cells into the peripheral regions. However, nor was there any indication of ectopic granulopoiesis. For this reason, the animal was diagnosed with granulocytic hyperplasia which could be considered a pre-myelogenous leukemic state.

#### AKR Mice at 100 ppm

The chronic exposure study of these mice to 100 ppm benzene ended after 505 days with the death of the last test animal. Treated mice received 337 exposures at an integrated mean benzene concentration of 100 ppm. There appeared to be no differences between test and control mice in median survival or in rate of weight gain (Table 1).

Hematological monitoring was suspended after 37 weeks due to advancing mortality in test mice. Statistically

significant lymphocytopenia was observed in treated mice vs. controls after one exposure week and this trend continued throughout the exposure. Lymphocyte depressions, however, were not as severe as those observed in the 300 ppm study with these animals (Table 6). Red cell levels of treated animals were depressed relative to controls and these depressions were statistically significant ( $\pm 2S_x$ ) for 9/20 monitorings (Table 6). Test mice showed a trend to granulocytosis throughout the exposure, however granulocyte levels were elevated by statistically significant amounts in only 3/20 monitorings.

Histological evaluation revealed few differences between test and control animals. Malignant lymphoma was confirmed in 29/49 treated animals and in 24/50 controls. The average lifespan of treated animals dying with malignant lymphoma was 247 days. The average lifespan for controls was 226 days. Bone marrow hypoplasia was found in 11 treated mice (22%) but in only one control. This compares with an 81% incidence of bone marrow hypoplasia found in these animals at 300 ppm.

#### Sprague-Dawley Rats at 100 ppm

The chronic exposure of these rats at 100 ppm is in progress with seventeen treated animals and nine controls still alive. Treated animals have received 428 exposures over 638 days at an integrated mean benzene concentration of 100 ppm.

Peripheral blood data indicates a trend in treated animals toward anemia and lymphocytopenia, however, the differences between test and control animals in red cell and lymphocyte levels has been rarely statistically significant ( $\pm 2S_x$ ) (Table 7).

Histological evaluation of rats dying during the study has commenced. One treated rat has died with a confirmed chronic myelogenous leukemia. Details of the histological and hematological evaluation are as follows. The animal died after 233 exposures to 100 ppm benzene and showed a progressively increasing white count for several months before succumbing. The white count eventually rose to greater than  $300,000/\text{mm}^3$  (control= $10,000/\text{mm}^3$ ). The peripheral smears showed marked increases in granulocytes shifted to the left. There was severe splenomegaly accompanied by an infarct which was visible at necropsy. The bone marrow was green and contained unsegmented granulocytes infiltrating into the periosteal regions. The liver, spleen, lymph nodes, and kidneys were perfused with myelopoietic elements. Death was ascribed to chronic myelogenous leukemia.

With the development of this case of chronic myelogenous leukemia, all treated and control rats have been screened monthly for hematologic abnormalities. To date, no control animals have presented signs consistent with leukemia. A second test rat has shown signs which may be consistent with the early stages of chronic myelogenous leukemia. White

counts have reached in excess of  $100,000/\text{mm}^3$  and a shift to the left has been observed. This animal now undergoes weekly hematologic monitoring.

Area B

Rats at 300 ppm

Because the red cell count data were equivocal for these animals at 300 ppm and because circulating cell counts are not necessarily good indicators of bone marrow toxicity, several additional red cell parameters were investigated at various times during the course of the chronic exposure.

Lactic dehydrogenase levels were monitored periodically from the twentieth week of exposure. Elevated levels were found in the test animals when compared with controls and the higher levels were significant ( $\pm 2S_x$ ) for assays performed on the twenty-first, twenty-second and twenty-seventh weeks (Table 8). Assays performed subsequent to the twenty-seventh week still showed elevated levels for test animals, however, control animals also showed increased LDH levels.

Red cell osmotic fragility was determined during the eighteenth, twenty-ninth and thirty-first weeks of exposure. Initially test animals showed slightly increased fragility at 0.35% saline. Subsequent determinations, however, showed no significant difference between test and control animals.

Red cell glutathione levels assayed on the fifteenth and seventeenth week of exposure showed no significant

differences between test and control animals nor did red cell acetylcholinesterase levels monitored during the twenty-seventh and twenty-eighth week of exposure.

AKR Mice at 300 ppm and 100 ppm

Since lactic acid dehydrogenase levels were found to be elevated at times during the exposure of Sprague-Dawley rats to 300 ppm benzene, it was decided to modify L.D.H. procedures so that this parameter could be followed during the exposure of AKR mice to 300 ppm and 100 ppm benzene vapor. No significant differences were found in the L.D.H. levels of test and control animals at either exposure level. Red cell glutathione levels were lower in test mice exposed to 300 ppm, however this decrease seemed to parallel the red blood cell drop found in test animals and therefore may not have been due to decreased production of glutathione.

Area C

AKR Mice at 300 ppm and 100 ppm

Cytogenetic analyses were performed on femoral cells from AKR mice exposed to 300 ppm benzene and from air controls. Assays were taken regularly until the ninety-third exposure day. The results are summarized in Table 9.

The mitotic rate was lower in benzene-treated mice except for the last three data points. The number of euploid cells was higher in the controls at each point in the study.



In the latter half of the study after five weeks of exposure, there were more breaks observed in treated than in untreated animals. Although the direction of change was the same, a statistically significant increment within the animal pairs was only seen at 38 and 93 days ( $p < 0.05$ ). Almost all breaks were open chromatid or chromosome breaks with the distal piece in geographic proximity to the parent chromosome, but with distal and lateral displacement. The breaks essentially followed the expected distribution for Poisson events with well over 90% of the cells in each series having only a single break.

A successful bone marrow cloning technique was developed using C-57 B1 and AKR mouse marrow cells. This procedure was applied on a regular basis to femoral marrow cells taken from AKR mice exposed to 100 ppm benzene. Although an initial depression in the number of colonies formed from benzene treated mice was noted, subsequent assays showed no differences in colony forming ability or cellularity (Table 10).

### Phase III

Techniques were developed during the course of these studies for the determination of benzene in blood and tissue.<sup>7,8</sup> Application of these techniques demonstrated that benzene concentrates in the bone marrow during a single 6 hr exposure (Table 11). In addition, the results indicated a correlation

between various tissue concentrations and atmospheric concentrations.

Pharmacokinetic analyses were performed on the time-courses of benzene in the blood of animals exposed to benzene vapor. Because of the extreme differences in hematological responses elicited from Sprague-Dawley rats and AKR mice exposed to given levels of benzene, these animals were chosen as models for this study. Animals were given 20 exposures to either 300 ppm or 100 ppm benzene and pharmacokinetic analyses were performed after the first, sixth and twentieth exposures. Results show that the mice eliminated benzene faster than the rats for a given atmospheric concentration (Table 12). In addition, mice exposed to 300 ppm showed a shift from a monoexponential clearance to a biexponential clearance after the twentieth exposure at 300 ppm. Such a shift is theoretically possible but does not seem to have been observed previously. Mice also showed a trend to increasing rates of elimination at 300 ppm which is indicative of metabolic enzyme induction.

The comparative rates of elimination for these rats and mice correlates well with the observed hematotoxicity. AKR mice exhibited a much more severe toxic response to a given level of benzene than the rats and the rates of elimination for the mice were faster than the rates observed for rats at both levels of benzene studied. In addition, at 300 ppm, AKR mice showed a continuing drop in peripheral cell levels

up to and beyond 20 exposures and this correlates well with the increasing rates of elimination observed for these mice at 300 ppm.

#### Summary

1. Cytopenic effects of benzene exposure were readily produced in the animals chosen for study at 300 ppm. At this level, all animals exhibited peripheral lymphocytopenia and all of the mice studied exhibited anemia. Rats, on the other hand, showed lymphocytopenia but only a trend toward anemia. At 100 ppm AKR mice gave evidence of a dose-response effect in that the lymphocytopenia and anemia were not as marked as at the 300 ppm level. Rats at 100 ppm also showed a dose-response cytopenic effect by exhibiting only non-statistically significant trends to peripheral cell effects.

2. The pharmacokinetic data shows a direct relationship between the rate of clearance of benzene and benzene-induced cytopenia. This situation is the reverse of the usual relationship observed between toxicity and clearance rates. For most toxic compounds, a faster clearance rate usually causes a decrease in administered dose and therefore milder toxicity. Our toxicity data suggest that the faster clearance rates exhibited by the mice vis a vis the rats may not be a function of increased respiration rates alone but may also be due to increased metabolic rates which would lead to greater accumulation of toxic metabolites. This view is

further supported by the fact that the clearance rates and cytotoxicity for AKR mice at 300 ppm appear to increase with increasing exposure. It would be difficult to ascribe these increasing rates to increased respiration rates. A more reasonable explanation would seem to be metabolic enzyme induction with consequent increases in toxic metabolites.

3. All of the mouse strains studied responded to the exposures with granulocytoses. At times, test mice showed six-fold increases in granulocyte levels vis a vis controls. Often shifts to immature cell types accompanied these increased granulocyte levels. For the CD-1 mice, a spectrum of various stages of granulocyte proliferation was noted. In view of the observed cytopenic effects of benzene on the lymphoid and erythroid cell lines, the observed proliferative effect on the myeloid cell line seems all the more compelling. This is especially true in view of the correlation between benzene exposure and myelogenous leukemia in humans.

4. There were three confirmed cases of myelogenous leukemia among the test animals studied. In view of the small numbers of animals developing leukemia, these results must be considered preliminary. Although the number of controls used in these studies was relatively small, there have been, to our knowledge, no reports of spontaneous myelogenous leukemia in either CD-1 mice or Sprague-Dawley rats. These three cases of myelogenous leukemia, therefore, should be considered indicative of a causal relationship.

between benzene exposure and myelogenous leukemia. Additional evidence for the carcinogenicity of benzene comes from the statistically significant increase in lymphoma among the C-57 Bl mice. Several known chemical carcinogens<sup>6</sup> as well as ionizing radiation produce a similar response in these mice.

5. The question of whether myelogenous leukemia follows episodes of peripheral cytopenia or arises without noticeable "preleukemic" peripheral cell changes must be considered as unanswered. The 2 mice developed leukemia after long episodes of peripheral anemia and lymphocytopenia whereas the rat showed peripheral cell counts within normal limits before presenting signs of myelogenous leukemia. It is possible that these varying responses represent species variability to a given toxic agent or they may have been due to varying individual responses. It is obvious that additional work should be conducted in this area.

References

1. Benzene Toxicity a critical evaluation. Sidney Laskin and Bernard D. Goldstein, Eds. J. Tox. and Environ. Health., Supplement 2, November, 1977.
2. Albrecht, A.M., Biedler, J.L., Hutchison, D.J., Spengler, B.A. and Stockert, E. Radiation-induced murine leukemic ERLD in cell culture. Cancer Research, 36:3784-3788, 1976.
3. Haas, M. Transient virus expression during murine leukemia induction by x-irradiation. J. Natl. Cancer Inst., 58:251-257, 1977.
4. Kaplan, H.S. On the natural history of murine leukemias. Cancer Research, 27:1325-1340, 1967.
5. Percy, D.H. and Jonas, A.M. Incidence of spontaneous tumors in CD-1 Ham/ICR mice. J. Natl. Cancer Inst., 46: 1045-1065, 1971.
6. Imamura, M., Nakano, M., Kawase, A., Kawamura, Y., and Yokoro, K. Synergistic action of N-Nitrosobutylurea and azathioprine in induction of leukemia in C-57 Bl mice. GANN, 64:493-498, 1973.
7. Snyder, C.A., Laskin, S. and Goldstein, B.D. An extractive method for the determination of benzene in blood. Am. Ind. Hyg. Assoc. J., 833-836, Nov., 1975.
8. Snyder, C.A., Erlichman, M.N., Goldstein, B.D., and Laskin, S. An extractive method to determine benzene in tissue by gas chromatography. Am. Ind. Hyg. Assoc. J., 38:272-276, June, 1977.

Table 1

<u>Animal</u>	<u>Number at Risk</u>	<u>Exposure Level (ppm)</u>	<u>Calculated* Median Survival Time</u>
Sprague-Dawley Rat	45	300	59 weeks
	27	0	73 weeks
Sprague-Dawley Rat	40	100	In progress
	40	0	.
AKR Mouse	60	300	17 weeks
	60	0	47 weeks
AKR Mouse	49	100	39 weeks
	50	0	41 weeks
CD-1 Mouse	40	300	31 weeks
	40	0	50 weeks
C-57 B1 Mouse	40	300	49 weeks
	40	0	83 weeks

\* The 50% survival point from least squares linear regression curve of log of survival vs. probability of mortality.

TABLE 2

COMPARISON OF RED BLOOD COUNTS BETWEEN SPRAGUE-DAWLEY RATS  
EXPOSED TO 300 PPM BENZENE AND AIR CONTROLS

Weeks After First Exposure	Test $\pm 2S_x \times 10^6$	Control $\pm 2S_x \times 10^6$
4	16.00 $\pm$ 1.70	17.20 $\pm$ 2.90
8	9.83 $\pm$ 1.20	9.60 $\pm$ 0.80
12	8.35 $\pm$ 0.68	8.11 $\pm$ 0.62
16	3.51 $\pm$ 0.46	4.61 $\pm$ 0.66
20	7.88 $\pm$ 0.60	8.60 $\pm$ 0.32
24	10.11 $\pm$ 1.16	8.77 $\pm$ 0.76
28	7.83 $\pm$ 0.34	8.55 $\pm$ 0.74
32	8.47 $\pm$ 0.30	8.74 $\pm$ 0.32
35	7.61 $\pm$ 0.04	9.00 $\pm$ 0.46
40	9.28 $\pm$ 0.26	9.20 $\pm$ 0.10
44	10.25 $\pm$ 1.06	9.59 $\pm$ 0.40
48	9.64 $\pm$ 0.52	9.70 $\pm$ 0.30
52	8.37 $\pm$ 0.86	9.01 $\pm$ 0.38



TABLE 3

WEIGHT CHANGES OF A/R MICE EXPOSED TO 300 PPM

BENZENE AND AIR CONTROLS

Weeks After First Exposure	Percent Weight Change*		Percent Weight Change* Air Controls Initial Weight = 28.4 g
	Exposed Mice Initial Weight = 28.4 g		
4	91		103
8	79		110
12	68		105
16	59		102
20	61		106
24	74		91
28	--		97
36	--		98
40	--		91
48	--		90
52	--		87

\* Initial Weight Taken as 100%

TABLE 4  
RETICULOCYTE COUNTS OF AKR MICE  
EXPOSED TO 300 PPM BENZENE AND AIR CONTROLS

Days After 1st Exposure Day	Benzene % Retic $\pm 2S_x$	Control % Retic $\pm 2S_x$
0	7.4 $\pm$ 2.0	5.1 $\pm$ 1.8
9	6.0 $\pm$ 1.2	5.2 $\pm$ 1.8
23	2.2 $\pm$ 0.8	1.6 $\pm$ 0.4
37	3.3 $\pm$ 0.8	1.5 $\pm$ 0.2
65	3.2 $\pm$ 1.0	1.2 $\pm$ 0.2

Table 5  
Comparison of Peripheral Blood Counts Between  
C-57 31 Mice Exposed to 300 ppm and Air Control

Weeks After First Exposure	Test			Control		
	<u>Lymphs</u>	<u>Polys</u>	<u>RBC</u>	<u>Lymphs</u>	<u>Polys</u>	<u>RBC</u>
0	9100	1000	10.67	7600	1000	10.79
1	6600*	1200*	9.43*	11900	1900	10.64
9	4000*	3300	6.19*	12000	1800*	9.39
17	2300*	4100	6.63*	11500	2300	9.88
25	2000*	4800	6.95*	10000	2600*	9.65
33	3200*	3800	6.48*	10500	2000*	9.48
41	4600*	4500	6.29*	11800	2500	9.66
49	3800*	8300	6.26*	12100	2000*	9.57
57	2500*	7200	6.59*	14800	3200*	9.78

\* denotes lower by a statistically significant difference ( $\pm$  2 standard errors).

Table 6

Comparison of Peripheral Cell Counts Between  
AKR Mice Exposed to 100 ppm and Air Controls

Weeks After First Exposure	Test			Control		
	<u>Lymphs</u>	<u>Polys</u>	<u>RBC</u>	<u>Lymphs</u>	<u>Polys</u>	<u>RBC</u>
0	10400	3300	9.49	9400	2600	9.33
1	6700*	1700*	7.93*	9800	2900	8.83
5	3600*	3900	8.40*	9300	4400	9.06
9	3200*	4000	8.26*	10900	3600	9.07
13	2700*	2800	8.12*	8000	3100	8.88
17	3500*	5000	8.80	9500	2600*	9.18
21	2800*	6000	8.54	7100	2500*	9.18
25	2400*	5000	8.27	6800	3800	9.03
29	4200	4000	8.21	4500	3600	10.17
33	2800*	5100	8.56	7600	4300	9.82
37	2500*	7000	6.90*	7800	5100	8.14

\* denotes lower by a statistically significant difference ( $\pm$  2 standard errors).

Table 7  
Comparison of Peripheral Blood Counts Between  
Sprague-Dawley Rats Exposed to 100 ppm and Air Control

Weeks After First Exposure	Test			Control		
	<u>Lymphs</u>	<u>Polys</u>	<u>RbC</u>	<u>Lymphs</u>	<u>Polys</u>	<u>RBC</u>
0	14100	2500	6.59	13900	2900	6.98
8	12100	1400	7.50	11900	2400	7.57
16	9100	2000	7.70	11200	1900	8.09
24	9300	2200	7.67	10700	1300*	7.95
32	10600	2900	8.04	13800	3900	8.20
38	9900	2400	8.60	11400	2300	8.77
45	10300	2900	7.97	11900	3000	8.47
53	11300	3700	8.03	12700	4300	8.30
61	11200	3600*	7.68*	12000	6100	9.00
70	10621	5100	8.21	13200	5200	9.08
77	9000	4700	8.05	11800	5700	8.40

\* denotes lower by a statistically significant difference ( $\pm 2$  standard errors).

TABLE 8

COMPARISON OF LACTIC ACID DEHYDROGENASE LEVELS BETWEEN  
SPRAGUE-DAWLEY RATS EXPOSED TO 300 PPM BENZENE AND AIR CONTROLS

Weeks After First Exposure	Controls In L.D.H. Units (C)	2 S <sub>x</sub>	Exposed In L.D.H. Units (E)	2 S <sub>x</sub>	Differences In L.D.H. Units (E - C)
20	570	270	860	270	+ 290
21	650	130	1410	300	+ 760
22	1000	310	1530	200	+ 530
27	270	70	1040	350	+ 770
28	530	120	940	300	+ 410
36	1110	490	1150	350	+ 40

TABLE 9

COMPARISON OF BONE MARROW CELLULAR MORPHOLOGY BETWEEN AKR MICE  
EXPOSED TO 300 PPM BENZENE AND AIR CONTROLS

Exposure	Duration of Exposure (Days)	Percent Diploid	Percent Hypodiploid	Percent Hypodiploid	Percent Cells With Breaks	Mitotic Rates
Pre-Exposure	--	90	0	1	5	1.8
Benzene	10	74	26	0	21	0.6
Air	--	91	7	2	16	0.8
Benzene	17	31	19	0	20	1.2
Air	--	86	13	1	21	2.0
Benzene	24	86	11	3	21	1.0
Air	--	93	7	0	30	2.0
Benzene	38	79	18	3	29	1.6
Air	--	96	4	0	16	2.0
Benzene	52	80	20	0	12	1.0
Air	--	92	5	3	8	1.0
Benzene	66	86	6	8	26	1.6
Air	--	92	8	0	21	1.2
Benzene	93	85	15	0	34	3.0
Air	--	91	9	0	16	1.6

TABLE 10  
COMPARISON OF CELLULAR PROLIFERATIVE ABILITY OF BONE MARROW CELLS  
FROM AKR MICE EXPOSED TO 100 PPM BENZENE AND AIR CONTROLS

Weeks After First Exposure	Type of Exposure	Number of Colonies	Cellularity x 10 <sup>6</sup>
2	Benzene	1	15
	Benzene	5	19
	Air	35	28
	Air	21	20
6	Benzene	13	34
	Benzene	13	29
	Air	19	44
	Air	15	35
10	Benzene	2	28
	Benzene	2	30
	Air	2	30
	Air	4	20
12	Benzene	7	16
	Benzene	5	20
	Air	1	39
	Air	12	25



TABLE 11  
BENZENE ( $C_6H_6$ ) DISTRIBUTION IN HEMATOPOIETIC TISSUE OF AKR MICE  
EXPOSED TO BENZENE VAPOR FOR 6 HOURS

100 PPM

Mouse	Weight (g)	$C_6H_6$ Conc. in blood ( $\mu g/ml$ )	$C_6H_6$ Conc. in liver ( $\mu g/g$ )	$C_6H_6$ Conc. in spleen ( $\mu g/g$ )	$C_6H_6$ Conc. in marrow ( $\mu g/g$ )
1	27	2.22	17.58	2.50	3.58
2	33	3.21	7.10	1.55	5.85
3	30	8.47	13.67	2.07	1.71
4	36	1.81	10.55	0.43	24.34

300 PPM

5	29	13.98	53.69	2.38	11.40
6	29	11.51	90.41	6.70	41.35
7	34	9.29	57.00	19.61*	15.63
8	28	23.07	58.25	11.17	75.66

\* Splenomegaly

Table 12

Comparison of Mean Elimination Rate Constants  
Between Rats and Mice Exposed to Benzene Vapor

Concentration	Exposure		
	1st	6th	20th
300 ppm	$k_m/k_r = 4.3$	$k_m/k_r = 4.2$	$k_m/k_r = 7.0$
100 ppm	$k_m/k_r = 5.3$	$k_m/k_r = 3.4$	$k_m/k_r = 5.2$

$k_m$  = mean rate constants for mice in reciprocal minutes

$k_r$  = mean rate constants for rats in reciprocal minutes

Test Lymph  
Control Lymph  
Test Polys.  
Control Polys.

Figure 1 S-D Rats at 300ppm






Figure 2  
AKR Mice at 300ppm



# RBC of AKR Mice at 300 ppm

Figure 3







Figure 5





UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

Barbara J. Price  
Vice President  
Health, Environment & Safety  
Phillips Petroleum Company  
Bartlesville, Oklahoma 74004

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

MAY 08 1995

EPA acknowledges the receipt of information submitted by your organization under Section 8(e) of the Toxic Substances Control Act (TSCA). For your reference, copies of the first page(s) of your submission(s) are enclosed and display the TSCA §8(e) Document Control Number (e.g., 8EHQ-00-0000) assigned by EPA to your submission(s). Please cite the assigned 8(e) number when submitting follow-up or supplemental information and refer to the reverse side of this page for "EPA Information Requests".

All TSCA 8(e) submissions are placed in the public files unless confidentiality is claimed according to the procedures outlined in Part X of EPA's TSCA §8(e) policy statement (43 FR 11110, March 16, 1978). Confidential submissions received pursuant to the TSCA §8(e) Compliance Audit Program (CAP) should already contain information supporting confidentiality claims. This information is required and should be submitted if not done so previously. To substantiate claims, submit responses to the questions in the enclosure "Support Information for Confidentiality Claims". This same enclosure is used to support confidentiality claims for non-CAP submissions.

Please address any further correspondence with the Agency related to this TSCA 8(e) submission to:

Document Processing Center (7407)  
Attn: TSCA Section 8(e) Coordinator  
Office of Pollution Prevention and Toxics  
U.S. Environmental Protection Agency  
Washington, D.C. 20460-0001

EPA looks forward to continued cooperation with your organization in its ongoing efforts to evaluate and manage potential risks posed by chemicals to health and the environment.

Sincerely,

*Terry R. O'Bryan*  
Terry R. O'Bryan  
Risk Analysis Branch

Enclosure

12578A



Recycled/Recyclable  
Printed with Soy/Canola Ink on paper that  
contains at least 50% recycled fiber



H

### Triage of 8(e) Submissions

Date sent to triage: 12/14/95

NON-CAP

CAP

Submission number: 12578A

TSCA Inventory:

Y

N

D

Study type (circle appropriate):

Group 1 - Dick Clements (1 copy total)

ECO

AQUATO

Group 2 - Ernie Falke (1 copy total)

ATOX

SBTOX

SEN

w/NEUR

Group 3 - Elizabeth Margosches (1 copy each)

STOX

CTOX

EPI

RTOX

GTOX

STOX/ONCO

CTOX/ONCO

IMMUNO

CYTO

NEUR

Other (FATE, EXPO, MET, etc.):

Notes:

**THIS IS THE ORIGINAL 8(e) SUBMISSION; PLEASE REFILE AFTER TRIAGE DATABASE ENTRY**

#### For Contractor Use Only

entire document: 0 1 2 pages 1 pages 4, 2, tabs

Notes:

Contractor reviewer :

LPS

Date:

4/14/95

CECATS DATA: 0992-12578 SEQ. A

Submission # SEHO

TYPE: INT. SUPP FLWP

SUBMITTER NAME:

Phillips Petroleum

Company

INFORMATION REQUESTED: FLWP DATE:

0201 NO INFO REQUESTED

0202 INFO REQUESTED (TECH)

0203 INFO REQUESTED (VOL. ACTIONS)

0204 INFO REQUESTED (REPORTING RATIONALE)

DISPOSITION:

0205 REFER TO CHEMICAL SCREENING

0206 CAP NOTICE

VALUATION ACTIONS:

0201 NOT AT RISK FOR ITD

0202 STUDIES PLANNED/IN PROGRESS

0203 INTERACTION OF WORKING MATERIALS

0204 LABELS AND CHANGES

0205 PROCESSING AND LOGISTICS

0206 APPROPRIATE DISCONTINUED

0207 PRODUCTION DISCONTINUED

0208 CONFIDENTIAL

SUB. DATE: 08/24/92

OTS DATE: 09/02/92

CRAD DATE:

03/07/95

CHEMICAL NAME:

Benzene

CASE

71-43-2

## INFORMATION TYPE:

PFC

## INFORMATION TYPE:

PFC

## INFORMATION TYPE:

PFC

0201	ONCO (HUMAN)	01 02 04	0216	EPICLIN	01 02 04	0241	IMMUNO (ANIMAL)	01 02 04
0202	ONCO (ANIMAL)	01 02 04	0217	HUMAN EXPOS (PROD CONTAM)	01 02 04	0242	IMMUNO (HUMAN)	01 02 04
0203	CELL TRANS (IN VITRO)	01 02 04	0218	HUMAN EXPOS (ACCIDENTAL)	01 02 04	0243	CHEMOPHY PROP	01 02 04
0204	MULTA (IN VITRO)	01 02 04	0219	HUMAN EXPOS (MONITORING)	01 02 04	0244	CLASTO (IN VITRO)	01 02 04
0205	MULTA (IN VIVO)	01 02 04	0220	BIOAQUA TOX	01 02 04	0245	CLASTO (ANIMAL)	01 02 04
0206	REPRO/TERATO (HUMAN)	01 02 04	0221	ENV. OCCURRENCE/FATE	01 02 04	0246	CLASTO (HUMAN)	01 02 04
0207	REPRO/TERATO (ANIMAL)	01 02 04	0222	ENV. ENCL OF ENV CONTAM	01 02 04	0247	DNA DAMAGE/PAIR	01 02 04
0208	NEURO (HUMAN)	01 02 04	0223	RESPONSE REQS: DELAY	01 02 04	0248	PRODUSE/PROC	01 02 04
0209	NEURO (ANIMAL)	01 02 04	0224	PRODUSE/PROC: ID	01 02 04	0251	MSDS	01 02 04
0210	ACUTE TOX (HUMAN)	01 02 04	0225	REPORTING RATIONALE	01 02 04	0259	OTHER	01 02 04
0211	CHR. TOX (HUMAN)	01 02 04	0226	CONFIDENTIAL	01 02 04			
0212	ACUTE TOX (ANIMAL)	01 02 04	0227	ALLERG (HUMAN)	01 02 04			
0213	SUB ACUTE TOX (ANIMAL)	01 02 04	0228	ALLERG (ANIMAL)	01 02 04			
0214	SUB CHRONIC TOX (ANIMAL)	01 02 04	0229	METAPHARMACO (ANIMAL)	01 02 04			
0215	CHRONIC TOX (ANIMAL)	01 02 04	0230	METAPHARMACO (HUMAN)	01 02 04			

## IMAGE DATA: NON-CBI INVENTORY

## ONGOING REVIEW

## SPECIES

## TOXICOLOGICAL CONCERN

## USE:

## PRODUCTION:

YES

YES (DROP/REFER)

RAT

LOW

CAS SR

NO

NO (CONTINUE)

MUS

MED

IN TITRATION

RAT

HIGH

Ctox

UNTESTED Rats and mice were exposed by inhalation to benzene for 6 hours/day & 5d/week for 9 weeks at 100 & 300 ppm. Main adverse effect was on blood & hematopoietic tissues. These changes were more pronounced at 300 ppm exposure and included leukopenia, increased hemoglobin fragments of spleen and bone marrow hypoplasia, red cells glutathione along with reduced red blood cells. No histopathology provided.

CECATR DATE: 1092-12584 SEQ. A

TYPE: INT SUP FLWP

SUBMITTER NAME: EIR Abbeon North

Ameca, Inc.

INFORMATION REQUESTED: FLWP DATE: 09/11/92  
 601 NO INFO REQUESTED  
 602 INFO REQUESTED (TECH)  
 603 INFO REQUESTED (VOL. ACTIONS)  
 604 INFO REQUESTED (REPORTING RATIONALE)  
 605 INFO REQUESTED (REPORTING RATIONALE)  
 606 INFO REQUESTED (REPORTING RATIONALE)  
 607 INFO REQUESTED (REPORTING RATIONALE)  
 608 INFO REQUESTED (REPORTING RATIONALE)  
 609 INFO REQUESTED (REPORTING RATIONALE)  
 610 INFO REQUESTED (REPORTING RATIONALE)  
 611 INFO REQUESTED (REPORTING RATIONALE)  
 612 INFO REQUESTED (REPORTING RATIONALE)  
 613 INFO REQUESTED (REPORTING RATIONALE)  
 614 INFO REQUESTED (REPORTING RATIONALE)  
 615 INFO REQUESTED (REPORTING RATIONALE)  
 616 INFO REQUESTED (REPORTING RATIONALE)  
 617 INFO REQUESTED (REPORTING RATIONALE)  
 618 INFO REQUESTED (REPORTING RATIONALE)  
 619 INFO REQUESTED (REPORTING RATIONALE)  
 620 INFO REQUESTED (REPORTING RATIONALE)  
 621 INFO REQUESTED (REPORTING RATIONALE)  
 622 INFO REQUESTED (REPORTING RATIONALE)  
 623 INFO REQUESTED (REPORTING RATIONALE)  
 624 INFO REQUESTED (REPORTING RATIONALE)  
 625 INFO REQUESTED (REPORTING RATIONALE)  
 626 INFO REQUESTED (REPORTING RATIONALE)  
 627 INFO REQUESTED (REPORTING RATIONALE)  
 628 INFO REQUESTED (REPORTING RATIONALE)  
 629 INFO REQUESTED (REPORTING RATIONALE)  
 630 INFO REQUESTED (REPORTING RATIONALE)  
 631 INFO REQUESTED (REPORTING RATIONALE)  
 632 INFO REQUESTED (REPORTING RATIONALE)  
 633 INFO REQUESTED (REPORTING RATIONALE)  
 634 INFO REQUESTED (REPORTING RATIONALE)  
 635 INFO REQUESTED (REPORTING RATIONALE)  
 636 INFO REQUESTED (REPORTING RATIONALE)  
 637 INFO REQUESTED (REPORTING RATIONALE)  
 638 INFO REQUESTED (REPORTING RATIONALE)  
 639 INFO REQUESTED (REPORTING RATIONALE)  
 640 INFO REQUESTED (REPORTING RATIONALE)  
 641 INFO REQUESTED (REPORTING RATIONALE)  
 642 INFO REQUESTED (REPORTING RATIONALE)  
 643 INFO REQUESTED (REPORTING RATIONALE)  
 644 INFO REQUESTED (REPORTING RATIONALE)  
 645 INFO REQUESTED (REPORTING RATIONALE)  
 646 INFO REQUESTED (REPORTING RATIONALE)  
 647 INFO REQUESTED (REPORTING RATIONALE)  
 648 INFO REQUESTED (REPORTING RATIONALE)  
 649 INFO REQUESTED (REPORTING RATIONALE)  
 650 INFO REQUESTED (REPORTING RATIONALE)  
 651 INFO REQUESTED (REPORTING RATIONALE)  
 652 INFO REQUESTED (REPORTING RATIONALE)  
 653 INFO REQUESTED (REPORTING RATIONALE)  
 654 INFO REQUESTED (REPORTING RATIONALE)  
 655 INFO REQUESTED (REPORTING RATIONALE)  
 656 INFO REQUESTED (REPORTING RATIONALE)  
 657 INFO REQUESTED (REPORTING RATIONALE)  
 658 INFO REQUESTED (REPORTING RATIONALE)  
 659 INFO REQUESTED (REPORTING RATIONALE)  
 660 INFO REQUESTED (REPORTING RATIONALE)  
 661 INFO REQUESTED (REPORTING RATIONALE)  
 662 INFO REQUESTED (REPORTING RATIONALE)  
 663 INFO REQUESTED (REPORTING RATIONALE)  
 664 INFO REQUESTED (REPORTING RATIONALE)  
 665 INFO REQUESTED (REPORTING RATIONALE)  
 666 INFO REQUESTED (REPORTING RATIONALE)  
 667 INFO REQUESTED (REPORTING RATIONALE)  
 668 INFO REQUESTED (REPORTING RATIONALE)  
 669 INFO REQUESTED (REPORTING RATIONALE)  
 670 INFO REQUESTED (REPORTING RATIONALE)  
 671 INFO REQUESTED (REPORTING RATIONALE)  
 672 INFO REQUESTED (REPORTING RATIONALE)  
 673 INFO REQUESTED (REPORTING RATIONALE)  
 674 INFO REQUESTED (REPORTING RATIONALE)  
 675 INFO REQUESTED (REPORTING RATIONALE)  
 676 INFO REQUESTED (REPORTING RATIONALE)  
 677 INFO REQUESTED (REPORTING RATIONALE)  
 678 INFO REQUESTED (REPORTING RATIONALE)  
 679 INFO REQUESTED (REPORTING RATIONALE)  
 680 INFO REQUESTED (REPORTING RATIONALE)  
 681 INFO REQUESTED (REPORTING RATIONALE)  
 682 INFO REQUESTED (REPORTING RATIONALE)  
 683 INFO REQUESTED (REPORTING RATIONALE)  
 684 INFO REQUESTED (REPORTING RATIONALE)  
 685 INFO REQUESTED (REPORTING RATIONALE)  
 686 INFO REQUESTED (REPORTING RATIONALE)  
 687 INFO REQUESTED (REPORTING RATIONALE)  
 688 INFO REQUESTED (REPORTING RATIONALE)  
 689 INFO REQUESTED (REPORTING RATIONALE)  
 690 INFO REQUESTED (REPORTING RATIONALE)  
 691 INFO REQUESTED (REPORTING RATIONALE)  
 692 INFO REQUESTED (REPORTING RATIONALE)  
 693 INFO REQUESTED (REPORTING RATIONALE)  
 694 INFO REQUESTED (REPORTING RATIONALE)  
 695 INFO REQUESTED (REPORTING RATIONALE)  
 696 INFO REQUESTED (REPORTING RATIONALE)  
 697 INFO REQUESTED (REPORTING RATIONALE)  
 698 INFO REQUESTED (REPORTING RATIONALE)  
 699 INFO REQUESTED (REPORTING RATIONALE)  
 700 INFO REQUESTED (REPORTING RATIONALE)

VOLUNTARY ACTIONS:  
 601 NO ACTION IN PROGRESS  
 602 ACTION IN PROGRESS  
 603 ACTION IN PROGRESS  
 604 ACTION IN PROGRESS  
 605 ACTION IN PROGRESS  
 606 ACTION IN PROGRESS  
 607 ACTION IN PROGRESS  
 608 ACTION IN PROGRESS  
 609 ACTION IN PROGRESS  
 610 ACTION IN PROGRESS  
 611 ACTION IN PROGRESS  
 612 ACTION IN PROGRESS  
 613 ACTION IN PROGRESS  
 614 ACTION IN PROGRESS  
 615 ACTION IN PROGRESS  
 616 ACTION IN PROGRESS  
 617 ACTION IN PROGRESS  
 618 ACTION IN PROGRESS  
 619 ACTION IN PROGRESS  
 620 ACTION IN PROGRESS  
 621 ACTION IN PROGRESS  
 622 ACTION IN PROGRESS  
 623 ACTION IN PROGRESS  
 624 ACTION IN PROGRESS  
 625 ACTION IN PROGRESS  
 626 ACTION IN PROGRESS  
 627 ACTION IN PROGRESS  
 628 ACTION IN PROGRESS  
 629 ACTION IN PROGRESS  
 630 ACTION IN PROGRESS  
 631 ACTION IN PROGRESS  
 632 ACTION IN PROGRESS  
 633 ACTION IN PROGRESS  
 634 ACTION IN PROGRESS  
 635 ACTION IN PROGRESS  
 636 ACTION IN PROGRESS  
 637 ACTION IN PROGRESS  
 638 ACTION IN PROGRESS  
 639 ACTION IN PROGRESS  
 640 ACTION IN PROGRESS  
 641 ACTION IN PROGRESS  
 642 ACTION IN PROGRESS  
 643 ACTION IN PROGRESS  
 644 ACTION IN PROGRESS  
 645 ACTION IN PROGRESS  
 646 ACTION IN PROGRESS  
 647 ACTION IN PROGRESS  
 648 ACTION IN PROGRESS  
 649 ACTION IN PROGRESS  
 650 ACTION IN PROGRESS  
 651 ACTION IN PROGRESS  
 652 ACTION IN PROGRESS  
 653 ACTION IN PROGRESS  
 654 ACTION IN PROGRESS  
 655 ACTION IN PROGRESS  
 656 ACTION IN PROGRESS  
 657 ACTION IN PROGRESS  
 658 ACTION IN PROGRESS  
 659 ACTION IN PROGRESS  
 660 ACTION IN PROGRESS  
 661 ACTION IN PROGRESS  
 662 ACTION IN PROGRESS  
 663 ACTION IN PROGRESS  
 664 ACTION IN PROGRESS  
 665 ACTION IN PROGRESS  
 666 ACTION IN PROGRESS  
 667 ACTION IN PROGRESS  
 668 ACTION IN PROGRESS  
 669 ACTION IN PROGRESS  
 670 ACTION IN PROGRESS  
 671 ACTION IN PROGRESS  
 672 ACTION IN PROGRESS  
 673 ACTION IN PROGRESS  
 674 ACTION IN PROGRESS  
 675 ACTION IN PROGRESS  
 676 ACTION IN PROGRESS  
 677 ACTION IN PROGRESS  
 678 ACTION IN PROGRESS  
 679 ACTION IN PROGRESS  
 680 ACTION IN PROGRESS  
 681 ACTION IN PROGRESS  
 682 ACTION IN PROGRESS  
 683 ACTION IN PROGRESS  
 684 ACTION IN PROGRESS  
 685 ACTION IN PROGRESS  
 686 ACTION IN PROGRESS  
 687 ACTION IN PROGRESS  
 688 ACTION IN PROGRESS  
 689 ACTION IN PROGRESS  
 690 ACTION IN PROGRESS  
 691 ACTION IN PROGRESS  
 692 ACTION IN PROGRESS  
 693 ACTION IN PROGRESS  
 694 ACTION IN PROGRESS  
 695 ACTION IN PROGRESS  
 696 ACTION IN PROGRESS  
 697 ACTION IN PROGRESS  
 698 ACTION IN PROGRESS  
 699 ACTION IN PROGRESS  
 700 ACTION IN PROGRESS

SUB DATE: 09/11/92 ORB DATE: 10/04/92 CRAD DATE: 03/07/95

CHEMICAL NAME:

TRTO Bis(tri-n-butyltin)oxide 56-35-9

INFORMATION TYPE:

LEC

INFORMATION TYPE:

LEC

INFORMATION TYPE:

LEC

601 ONCO (HUMAN)  
 602 ONCO (ANIMAL)  
 603 CELL TRANS (IN VITRO)  
 604 MUTA (IN VITRO)  
 605 MUTA (IN VIVO)  
 606 REPRO/TERATO (HUMAN)  
 607 REPRO/TERATO (ANIMAL)  
 608 NEURO (HUMAN)  
 609 NEURO (ANIMAL)  
 610 ACUTE TOX. (HUMAN)  
 611 CHR. TOX. (HUMAN)  
 612 ACUTE TOX. (ANIMAL)  
 613 SUB ACUTE TOX. (ANIMAL)  
 614 SUB CHRONIC TOX. (ANIMAL)  
 615 CHRONIC TOX. (ANIMAL)

616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700

601 HUMAN EXPOS (PROD CONTAM)  
 602 HUMAN EXPOS (ACCIDENTAL)  
 603 HUMAN EXPOS (MONITORING)  
 604 HUMAN EXPOS (MONITORING)  
 605 HUMAN EXPOS (MONITORING)  
 606 HUMAN EXPOS (MONITORING)  
 607 HUMAN EXPOS (MONITORING)  
 608 HUMAN EXPOS (MONITORING)  
 609 HUMAN EXPOS (MONITORING)  
 610 HUMAN EXPOS (MONITORING)  
 611 HUMAN EXPOS (MONITORING)  
 612 HUMAN EXPOS (MONITORING)  
 613 HUMAN EXPOS (MONITORING)  
 614 HUMAN EXPOS (MONITORING)  
 615 HUMAN EXPOS (MONITORING)  
 616 HUMAN EXPOS (MONITORING)  
 617 HUMAN EXPOS (MONITORING)  
 618 HUMAN EXPOS (MONITORING)  
 619 HUMAN EXPOS (MONITORING)  
 620 HUMAN EXPOS (MONITORING)  
 621 HUMAN EXPOS (MONITORING)  
 622 HUMAN EXPOS (MONITORING)  
 623 HUMAN EXPOS (MONITORING)  
 624 HUMAN EXPOS (MONITORING)  
 625 HUMAN EXPOS (MONITORING)  
 626 HUMAN EXPOS (MONITORING)  
 627 HUMAN EXPOS (MONITORING)  
 628 HUMAN EXPOS (MONITORING)  
 629 HUMAN EXPOS (MONITORING)  
 630 HUMAN EXPOS (MONITORING)  
 631 HUMAN EXPOS (MONITORING)  
 632 HUMAN EXPOS (MONITORING)  
 633 HUMAN EXPOS (MONITORING)  
 634 HUMAN EXPOS (MONITORING)  
 635 HUMAN EXPOS (MONITORING)  
 636 HUMAN EXPOS (MONITORING)  
 637 HUMAN EXPOS (MONITORING)  
 638 HUMAN EXPOS (MONITORING)  
 639 HUMAN EXPOS (MONITORING)  
 640 HUMAN EXPOS (MONITORING)  
 641 HUMAN EXPOS (MONITORING)  
 642 HUMAN EXPOS (MONITORING)  
 643 HUMAN EXPOS (MONITORING)  
 644 HUMAN EXPOS (MONITORING)  
 645 HUMAN EXPOS (MONITORING)  
 646 HUMAN EXPOS (MONITORING)  
 647 HUMAN EXPOS (MONITORING)  
 648 HUMAN EXPOS (MONITORING)  
 649 HUMAN EXPOS (MONITORING)  
 650 HUMAN EXPOS (MONITORING)  
 651 HUMAN EXPOS (MONITORING)  
 652 HUMAN EXPOS (MONITORING)  
 653 HUMAN EXPOS (MONITORING)  
 654 HUMAN EXPOS (MONITORING)  
 655 HUMAN EXPOS (MONITORING)  
 656 HUMAN EXPOS (MONITORING)  
 657 HUMAN EXPOS (MONITORING)  
 658 HUMAN EXPOS (MONITORING)  
 659 HUMAN EXPOS (MONITORING)  
 660 HUMAN EXPOS (MONITORING)  
 661 HUMAN EXPOS (MONITORING)  
 662 HUMAN EXPOS (MONITORING)  
 663 HUMAN EXPOS (MONITORING)  
 664 HUMAN EXPOS (MONITORING)  
 665 HUMAN EXPOS (MONITORING)  
 666 HUMAN EXPOS (MONITORING)  
 667 HUMAN EXPOS (MONITORING)  
 668 HUMAN EXPOS (MONITORING)  
 669 HUMAN EXPOS (MONITORING)  
 670 HUMAN EXPOS (MONITORING)  
 671 HUMAN EXPOS (MONITORING)  
 672 HUMAN EXPOS (MONITORING)  
 673 HUMAN EXPOS (MONITORING)  
 674 HUMAN EXPOS (MONITORING)  
 675 HUMAN EXPOS (MONITORING)  
 676 HUMAN EXPOS (MONITORING)  
 677 HUMAN EXPOS (MONITORING)  
 678 HUMAN EXPOS (MONITORING)  
 679 HUMAN EXPOS (MONITORING)  
 680 HUMAN EXPOS (MONITORING)  
 681 HUMAN EXPOS (MONITORING)  
 682 HUMAN EXPOS (MONITORING)  
 683 HUMAN EXPOS (MONITORING)  
 684 HUMAN EXPOS (MONITORING)  
 685 HUMAN EXPOS (MONITORING)  
 686 HUMAN EXPOS (MONITORING)  
 687 HUMAN EXPOS (MONITORING)  
 688 HUMAN EXPOS (MONITORING)  
 689 HUMAN EXPOS (MONITORING)  
 690 HUMAN EXPOS (MONITORING)  
 691 HUMAN EXPOS (MONITORING)  
 692 HUMAN EXPOS (MONITORING)  
 693 HUMAN EXPOS (MONITORING)  
 694 HUMAN EXPOS (MONITORING)  
 695 HUMAN EXPOS (MONITORING)  
 696 HUMAN EXPOS (MONITORING)  
 697 HUMAN EXPOS (MONITORING)  
 698 HUMAN EXPOS (MONITORING)  
 699 HUMAN EXPOS (MONITORING)  
 700 HUMAN EXPOS (MONITORING)

601 ALLERG (HUMAN)  
 602 ALLERG (ANIMAL)  
 603 METABOLISM (HUMAN)  
 604 METABOLISM (ANIMAL)  
 605 METABOLISM (HUMAN)  
 606 METABOLISM (ANIMAL)  
 607 METABOLISM (HUMAN)  
 608 METABOLISM (ANIMAL)  
 609 METABOLISM (HUMAN)  
 610 METABOLISM (ANIMAL)  
 611 METABOLISM (HUMAN)  
 612 METABOLISM (ANIMAL)  
 613 METABOLISM (HUMAN)  
 614 METABOLISM (ANIMAL)  
 615 METABOLISM (HUMAN)  
 616 METABOLISM (ANIMAL)  
 617 METABOLISM (HUMAN)  
 618 METABOLISM (ANIMAL)  
 619 METABOLISM (HUMAN)  
 620 METABOLISM (ANIMAL)  
 621 METABOLISM (HUMAN)  
 622 METABOLISM (ANIMAL)  
 623 METABOLISM (HUMAN)  
 624 METABOLISM (ANIMAL)  
 625 METABOLISM (HUMAN)  
 626 METABOLISM (ANIMAL)  
 627 METABOLISM (HUMAN)  
 628 METABOLISM (ANIMAL)  
 629 METABOLISM (HUMAN)  
 630 METABOLISM (ANIMAL)  
 631 METABOLISM (HUMAN)  
 632 METABOLISM (ANIMAL)  
 633 METABOLISM (HUMAN)  
 634 METABOLISM (ANIMAL)  
 635 METABOLISM (HUMAN)  
 636 METABOLISM (ANIMAL)  
 637 METABOLISM (HUMAN)  
 638 METABOLISM (ANIMAL)  
 639 METABOLISM (HUMAN)  
 640 METABOLISM (ANIMAL)  
 641 METABOLISM (HUMAN)  
 642 METABOLISM (ANIMAL)  
 643 METABOLISM (HUMAN)  
 644 METABOLISM (ANIMAL)  
 645 METABOLISM (HUMAN)  
 646 METABOLISM (ANIMAL)  
 647 METABOLISM (HUMAN)  
 648 METABOLISM (ANIMAL)  
 649 METABOLISM (HUMAN)  
 650 METABOLISM (ANIMAL)  
 651 METABOLISM (HUMAN)  
 652 METABOLISM (ANIMAL)  
 653 METABOLISM (HUMAN)  
 654 METABOLISM (ANIMAL)  
 655 METABOLISM (HUMAN)  
 656 METABOLISM (ANIMAL)  
 657 METABOLISM (HUMAN)  
 658 METABOLISM (ANIMAL)  
 659 METABOLISM (HUMAN)  
 660 METABOLISM (ANIMAL)  
 661 METABOLISM (HUMAN)  
 662 METABOLISM (ANIMAL)  
 663 METABOLISM (HUMAN)  
 664 METABOLISM (ANIMAL)  
 665 METABOLISM (HUMAN)  
 666 METABOLISM (ANIMAL)  
 667 METABOLISM (HUMAN)  
 668 METABOLISM (ANIMAL)  
 669 METABOLISM (HUMAN)  
 670 METABOLISM (ANIMAL)  
 671 METABOLISM (HUMAN)  
 672 METABOLISM (ANIMAL)  
 673 METABOLISM (HUMAN)  
 674 METABOLISM (ANIMAL)  
 675 METABOLISM (HUMAN)  
 676 METABOLISM (ANIMAL)  
 677 METABOLISM (HUMAN)  
 678 METABOLISM (ANIMAL)  
 679 METABOLISM (HUMAN)  
 680 METABOLISM (ANIMAL)  
 681 METABOLISM (HUMAN)  
 682 METABOLISM (ANIMAL)  
 683 METABOLISM (HUMAN)  
 684 METABOLISM (ANIMAL)  
 685 METABOLISM (HUMAN)  
 686 METABOLISM (ANIMAL)  
 687 METABOLISM (HUMAN)  
 688 METABOLISM (ANIMAL)  
 689 METABOLISM (HUMAN)  
 690 METABOLISM (ANIMAL)  
 691 METABOLISM (HUMAN)  
 692 METABOLISM (ANIMAL)  
 693 METABOLISM (HUMAN)  
 694 METABOLISM (ANIMAL)  
 695 METABOLISM (HUMAN)  
 696 METABOLISM (ANIMAL)  
 697 METABOLISM (HUMAN)  
 698 METABOLISM (ANIMAL)  
 699 METABOLISM (HUMAN)  
 700 METABOLISM (ANIMAL)

601 BLOOD (ANIMAL)  
 602 BLOOD (HUMAN)  
 603 CHEMISTS PROF  
 604 CLASTO (IN VITRO)  
 605 CLASTO (ANIMAL)  
 606 CLASTO (HUMAN)  
 607 DNA DAMAGE/FAIR  
 608 PRODUCE/PROC  
 609 MADS  
 610 OTHER

601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700

INFORMATION: NON-CL INVENTORY

YES

YES (PROCESSED)

RAT

LOW

NEED

USE:

PRODUCTION:

CAS SR NO

NO (CONTINUE)

IN RANGE

RAT-8

HIGH

UNTESTED Rats (Wistar) received TRTO at dosing days 8, 0.5, 20, 80 and 320 mg/kg (3x/week) for 4 and 6 weeks. Treatment produced effects in organs including thymus, spleen, liver, thyroid and pituitary at 320 and 80 mg/kg. The morphologic changes included decreased leucocytes, lymphoid and hematopoietic. Microscopic lesions observed included thymic and spleen atrophy, multifocal necrosis of liver parenchyma, flattening of the epithelial lining of thyroid follicles and decreased TSH immunoreactive cells in pituitary.

The NOAEL was 20 mg/kg.

Table 9. Summary of functional assessment of specific and nonspecific resistance after short-term and long-term exposure in the rat

Dietary concentration (mg/kg)	6 weeks <sup>a</sup>		4-6 months		15-17 months	
	20	80	0.5	5	50	0.5
<u>Thymus-dependent immunity</u>						
<i>T. spiralis</i> infection: expulsion of adult worms	+	++				
resistance to muscle larvae	+	++	-	+	++	-
IgE response	+	+	-	+	++	-
Delayed type hypersensitivity to ovalbumin and tuberculin	+	++	-	-	-	-
IgM and IgG response to <i>T. spiralis</i> and ovalbumin	-	-	-	-	-	-
IgG response to sheep red blood cells	-	+				-
T-mitogen response, thymus	+	++	-	-	+	
T-mitogen response, spleen	-	+	-	-	-	-
<u>Nonspecific resistance</u>						
<i>L. monocytogenes</i> infection, splenic clearance	+	++	-	-	+	-
Natural cell-mediated cytotoxicity, peritoneal macrophages	+	+	-	-	-	
Natural cell-mediated cytotoxicity, splenic NK cells	-	+	-	-	+	+

<sup>a</sup> From Vos *et al.*, 1984.

- no suppression; + slight to moderate suppression; ++ strong suppression

CECATS/TRIAGE TRACKING DBASE ENTRY FORM

CECATS DATA: 0992-12578 SEQ. A

Submission # BEHQ

TYPE: INT. SUPP FLWP

SUBMITTER NAME: Phillips Petroleum

Company

INFORMATION REQUESTED: FLWP DATE

0501 NO INFO REQUESTED

0502 INFO REQUESTED (TECH)

0503 INFO REQUESTED (VOL ACTIONS)

0504 INFO REQUESTED (REPORTING RATIONAL P)

DISPOSITION:

0639 REFER TO CHEMICAL SCREENING

0678 CAP NOTICE

VOLUNTARY ACTIONS:

0401 NO ACTION REPORTED

0402 STUDIES PLANNED/IN PROGRESS

0403 NOTIFICATION OF WORKING CONDITIONS

0404 LABEL/MSDS CHANGES

0405 PROCESS/HANDLING CHANGES

0406 APPAUSE DISCONTINUED

0407 PRODUCTION DISCONTINUED

0408 CONFIDENTIAL

SUB. DATE: 08/24/92 OTS DATE: 09/02/92 CSRAD DATE: 03/07/95

CHEMICAL NAME:

CASE

71-43-2

INFORMATION TYPE:	P.F.C.	INFORMATION TYPE:	P.F.C.
0201 ONCO (HUMAN)	01 02 04	0241 IMMUNO (ANIMAL)	01 02 04
0202 ONCO (ANIMAL)	01 02 04	0242 IMMUNO (HUMAN)	01 02 04
0203 CELL TRANS (IN VITRO)	01 02 04	0243 CHEM/PHYS PROP	01 02 04
0204 MUTA (IN VITRO)	01 02 04	0244 CLASTO (IN VITRO)	01 02 04
0205 MUTA (IN VIVO)	01 02 04	0245 CLASTO (ANIMAL)	01 02 04
0206 REPRO/TERATO (HUMAN)	01 02 04	0246 CLASTO (HUMAN)	01 02 04
0207 REPRO/TERATO (ANIMAL)	01 02 04	0247 DNA DAM/REPAIR	01 02 04
0208 NEURO (HUMAN)	01 02 04	0248 PRODUCE/PROC	01 02 04
0209 NEURO (ANIMAL)	01 02 04	0251 MSDS	01 02 04
0210 ACUTE TOX. (HUMAN)	01 02 04	0299 OTHER	01 02 04
0211 CHR. TOX. (HUMAN)	01 02 04		
0212 ACUTE TOX. (ANIMAL)	01 02 04		
0213 SUB ACUTE TOX (ANIMAL)	01 02 04		
0214 SUB CHRONIC TOX (ANIMAL)	01 02 04		
0215 CHRONIC TOX (ANIMAL)	01 02 04		

USE: PRODUCTION:

TOXICOLOGICAL CONCERN:

SPECIES

ONGOING REVIEW

TRIAGE DATA: NON-CBI INVENTORY

YES (DROP/REFER)

YES

NO (CONTINUE)

NO

REFR

IN NAME

RAT

LOW

MUS

MED

HIGH

ND

10/15/92

12578A

L/L/M/ND

Carcinogenicity in the Sprague Dawley rat is of low concern. Male rats were exposed via inhalation to 0 or 300 ppm benzene (27 controls and 45 exposed) for 6 hours/day, 5 days/week for life. Median survival time was decreased in the exposed rats. Hematological effects were: significant leukopenia due to selective decreases in lymphocyte levels, and slight anemia. Histopathology showed significantly increased incidence of hemosiderin pigments in the spleen and increased incidence of fatty changes of the bone marrow. There was no evidence of leukemia or lymphomas. An additional experiment was in progress involving 40 mice/group exposed to 0 or 100 ppm using the same exposure protocol as for 300 ppm. Histological analysis at 100 ppm was not complete, but one treated rat died due to chronic myelogenous leukemia. One treated rat that had not died also showed evidence of chronic myelogenous leukemia, but this leukemia had not been observed in any control rats. Although results at the low concentration were not complete, this study was rated low concern based on the absence of carcinogenic findings at the high concentration. However, it is unclear how complete the histopathological analysis was.

Inhalation carcinogenicity in the AKR mouse is of low concern. Groups of 60 male mice were exposed to 0 or 300 ppm for 6 hours/day, 5 days/week for life; separate testing was conducted with 50 mice/group exposed to 100 ppm and concurrent controls using the same exposure protocol. Survival time was markedly reduced at 300 ppm, but not at 100 ppm. No mice survived exposure to 300 ppm for more than 28 weeks, and severe weight loss was observed at this level. Lymphocytopenia and anemia were observed with concentration-related severity. The incidence of malignant lymphoma was lower at 300 ppm (2%) than in the control mice (91%), due to the high background rate in this strain and the median survival time in the exposed mice being shorter than the time to the first tumor in the controls. However, cytogenetic analysis showed increased aneuploidy of bone marrow cells from treated animals. At 100 ppm, the incidence of malignant lymphoma was comparable in the treated group (29/49) and the controls (24/50). Noncancer histological effects observed were increased incidences of hemosiderin pigments in the spleen and bone marrow hypoplasia at both treatment levels. This study is limited by the incomplete identification of the organs that underwent histopathological analysis.

Inhalation carcinogenicity in the C-57 B1 mouse is of medium concern. Groups of 40 male mice were exposed to 0 or 300 ppm for 6 hours/day, 5 days/week for life. Median survival time in exposed mice was 49 weeks, versus 83 weeks for the controls, and the exposed mice showed decreased weight gain compared to the controls for the duration of the study. Hematological findings in exposed mice were lymphocytopenia, anemia, granulocytosis, and hypoplastic bone

marrow. Malignant lymphoma was significantly increased in exposed mice (9/40) versus the controls (2/40). In addition, 13/31 exposed mice showed bone marrow hyperplasia and 15/31 had spleen hyperplasia, compared with no cases of bone marrow hyperplasia and 2/38 of spleen hyperplasia in the controls. This study is limited by the incomplete identification of the organs that underwent histopathological analysis.

Inhalation carcinogenicity in the CD-1 mouse is not determined, due to incomplete histological data for the treated group and no reported histological data for the control group. Groups of 40 male mice were exposed to 0 or 300 ppm for 6 hours/day, 5 days/week for life. Median survival time in exposed mice was about 60% that of the controls, and the exposed mice showed decreased weight gain compared to the controls for the duration of the study. Hematological findings in exposed mice were lymphocytopenia, anemia, and granulocytosis. Histological analysis was not complete, but histology was described for three treated mice, one dying of chronic myelogenous leukemia, one dying of acute myelogenous leukemia, and one diagnosed as having granulocytic hyperplasia, which was considered a pre-myelogenous leukemic state.